



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of it follows: Bill pinkish dusky toward tip (nail positively black) lamellated; trifle broader towards tip than at base; nostrils in the basal third. Head and about one-half neck brownish gray, somewhat mottled with white on sides and below. Under parts including linings of wings, lower wing coverts and axillars white. Upper parts generally, including upper wing coverts white. Primaries sooty; about a terminal fourth of the secondaries dark sooty brown, with a green speculum, each secondary slightly edged and tipped below with lighter. The white tail has all but two outer feathers dark tipped. Scapulars brown. A ring across the shoulders, part way round on the breast, chestnut. Feet flesh color.

Dimensions in inches: culmen (curve) 1.87; culmen (chord) 1.87; depth of bill 1; gape gape 2.12; length 22; wing 12.25; tail 4; mid-tow and claw 2.65; tarsus 2.65.—E. H. FORBUSH, 136 State House, Boston, Mass.

White Pelican (*Pelecanus erythrorhynchos*) Nesting in Minnesota.—From Dr. Frank M. Chapman's most excellent work, 'Camps and Cruises of an Ornithologist,' page 371, I quote the following: "In Western Minnesota, Pelicans nested as recently as 1878." While living at Aitkin, Minnesota, on August 15, 1904, a hunter brought to me two young Pelicans which he shot on the shores of Sandy Lake, about 40 miles north of Aitkin. They were small, not more than one-fourth grown, and were undoubtedly hatched and reared near the point of capture for they could not have migrated very far, in fact, were unable to fly.

Unfortunately their plumage was so badly soiled and so carelessly handled by the sportsman that I did not preserve them, which I have regretted since for the most undisputable evidence is the specimen in hand.—ALBERT LANO, Fayetteville, Ark.

American Egret (*Casmerodius egretta*) in Maine.—In 'The Auk' (Vol. XXXVIII, p. 109, January, 1921), I recorded an additional record of the Egret for the State of Maine—two specimens, one of which was shot and later secured by Dr. William P. Coues, who has since presented it to the Boston Society of Natural History. I also quoted twelve previously published records.

On July 15, 1921, Dr. William P. Coues saw a flock of seven on the Scarborough salt marshes which border a corner of the Prout's Neck Golf Links, and three days later a flock of nine were seen in the same locality by his brother, Robert W. Coues. This flock evidently scattered over the marshes, for between that date and the first week in September, only single birds or occasional pairs, were observed. Between July 19 and September 6, I observed single birds every few days. One favorite place was within 150 yards of the highway where there was constant passing. I stalked one bird and got within 75 yards, so that the yellow bill and black legs could plainly be seen with a glass. This bird was busily feeding in a ditch at low tide, but finally saw me and moved off to another part of the marsh.